

2008 12

100280

MATHEMATICS (Optional)**गणित (वैकल्पिक)****Time : 3 hours****Maximum Marks : 200****Note :**

- (i) In all attempt **Five** questions.
- (ii) Question No. 1 is **compulsory**.
- (iii) Of the remaining questions, Attempt **Any Four** by selecting **one question** from each section.
- (iv) Number of optional questions upto the prescribed number in the order in which questions have been solved, will only be assessed and excess answers of the question/s will not be assessed.
- (v) Candidate should not write roll number, any names (including his/her own), signature, address or any indication of his/her identity anywhere inside the answer book otherwise he/she will be penalised.

1. Answer any four of the following :

- (a) Define (i) Feasible solution to LPP. 2
(ii) Basic feasible solution to LPP. 2
(iii) Convex set. 2
- Show that the set of all feasible solution to a LPP (if a feasible solution exists) is a convex set. 4
- (b) If $\{a_n\}$ is a non-increasing sequence of positive numbers and if $\sum_{n=0}^{\infty} 2^n a_{2^n}$ converges, then show that $\sum_{n=0}^{\infty} a_n$ converges. 10
- (c) Determine components of velocity and acceleration of a moving particle along radial and transverse directions. 5 + 5
- (d) Write a programme in C/recent computer language to evaluate the roots of a quadratic equation $ax^2+bx+c = 0$ requesting the user to input the values of a, b, c and to output real roots root1 and root2. 10
- (e) If $\{f_n\}$ is a sequence of continuous real valued functions on the metric space X that converges uniformly to 'f' on X, then show that 'f' is also continuous on X. 10

P.T.O.

SECTION - A

2. (a) Show that every group is isomorphic to a subgroup of a permutation group $A(S)$ for some appropriate S . 18
State the name of this theorem. 2
- (b) If $\{v_1, v_2, \dots, v_n\}$ is a basis of a vector space V and if $\{w_1, w_2, \dots, w_m\}$ is linearly independent in V , then show that $m \leq n$. 20
3. (a) Define (i) Euclidean Domain. 3
(ii) Principal Ideal Domain (PID). 3
Show that every Euclidean Domain is PID. 14
- (b) State Cayley Hamilton theorem and using it find inverse of the matrix A if it exists. 2

$$A = \begin{pmatrix} 1 & 2 & 3 \\ 2 & 2 & 4 \\ 3 & 4 & 8 \end{pmatrix} \quad 18$$

SECTION - B

4. (a) If ' f ' is a real-valued continuous function on a compact metric space X , then show that $f(X)$, range of ' f ' is compact and ' f ' attains a maximum and minimum at points of X . 10 + 10
- (b) Define absolute convergence and conditional convergence for improper integrals of the type $\int_a^{\infty} f(x) dx$ for continuous function $f(x)$. 2 + 2
- Show that $\int_a^{\infty} \frac{\sin x}{x} dx$ is convergent but not absolutely. 8 + 8
5. (a) Define differentiability of a function of two variables at a point. 2
Let $f: E \rightarrow \mathbb{R}$ be defined on a neighbourhood E of $(a,b) \in \mathbb{R} \times \mathbb{R}$ such that $\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}$ are continuous at (a,b) . Show that ' f ' is differentiable at (a, b) . Is the converse of this is true? Justify your answer. 10
8

- (b) Define a Riemann integral for a bounded real function on $[a, b]$. 6
 Show that a bounded real function ' f ' is Riemann integrable on $[a, b]$ if and only if for every $\epsilon > 0$, there is a partition P of $[a, b]$ such that $U(P, f) - L(P, f) < \epsilon$. 7 + 7

SECTION - C

6. (a) Show that the general equation $ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$ of second degree represents a conic section. 20
 (b) Define (i) Divergence of a vector point function. 2
 (ii) Curl of a vector point function. 2
 (c) Show that $\bar{b} \cdot \nabla(\bar{a} \cdot \nabla \frac{1}{r}) = \frac{3(\bar{a} \cdot \bar{r})(\bar{b} \cdot \bar{r})}{r^5} - \frac{\bar{a} \cdot \bar{b}}{r^3}$ 10
 (d) Show that $\text{curl}(\bar{r} \times \bar{a}) = -2\bar{a}$. 6
7. (a) (i) Find the length of perpendicular from a point (x_1, y_1, z_1) to the plane $ax + by + cz = d$. 10
 (ii) Find equation of a sphere for which the circle $x^2 + y^2 + z^2 + 7y - 2z + 2 = 0$, $2x + 3y + 4z = 8$ is a great circle. 10
 (b) Verify Gauss divergence theorem for $\vec{f} \equiv (x^2 - yz)i + (y^2 - zx)j + (z^2 - xy)k$ taken over the rectangular parallelepiped $0 \leq x \leq a$, $0 \leq y \leq b$, $0 \leq z \leq c$. 20

SECTION - D

8. (a) (i) Explain when $M(x, y) + N(x, y)y' = 0$ is said to be exact in some rectangle R . 2
 (ii) Define integrating factor of the equation $M(x, y)dx + N(x, y)dy = 0$, where M and N have continuous partial derivatives in some rectangle R . 2
 Show that the equation $M(x, y) + N(x, y)y' = 0$ is exact in
 $R : |x - x_0| \leq a, |y - y_0| \leq b$ if and only if $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$ in R . 8 + 8
 (b) Find the inverse of the coefficient matrix of the system.

$$\begin{bmatrix} 1 & 1 & 1 \\ 4 & 3 & -1 \\ 3 & 5 & 3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 1 \\ 6 \\ 4 \end{bmatrix}$$

by Gauss - Jordan method with partial pivoting and hence solve the system. 20

P.T.O.

9. (a) If ϕ_1 is a solution of $y'' + a_1(x)y' + a_2(x)y = 0$ on an interval I, and $\phi_1(x) \neq 0$ on I, describe a method to determine a second linearly independent solution ϕ_2 of this differential equation on I. 12

Hence or otherwise find second linearly independent solution of $y'' - 4xy' + (4x^2 - 2)y = 0$ after verifying that $\phi_1(x) = e^{x^2}$ is a solution of this differential equation. 8

- (b) Describe Trapezoidal and Simpson $\frac{1}{3}rd$ rule to find $\int_a^b f(x)dx$ numerically. 5 + 5

Find an approximate value of $\int_0^1 \frac{dx}{1+x}$ by using Trapezoidal and Simpson $\frac{1}{3}rd$

rule and compare with exact solution. 5 + 5

- o O o -